MISSOURI BOTANICAL GARDEN
PETER H. RAVEN LIBRARY



GEORGE ENGELMANN BOTANICAL NOTEBOOKS

Pagination Note:
Since many of the items lack a specific page number, the page number displayed online refers to the sequentially created number each item was given upon cataloging the materials.

No. New York 185

Continental Bank

Pay to or order

Dollars

$

No. New York 185

Continental Bank

Pay to or order

Dollars

$

No. New York 185

Continental Bank

Pay to

Schott

A. Schott's Cactaceæ.

Drawings 1. Opuntia fragilis El Paso

2. O. Longispina, purple joints

3. O. pusilla see Remarks page 5. Tunita is the fruit

4 O. arborescens Remarks pg 6 Coyonoste is the plant

5. E. setispinus on Cretaceous rocks — see also living specimen

6. E. pulchellus Cayle pass, 13 ribs.

7. Cereus tabularis gracilis new Mex — no spines! & Chavon hills 3 miles ledge lands and ridges — see letter March 29th & Jan 13th — № 1. (Octo 5th 1853) also

8 Cereus (Pilella) Organos, Cucumber plant, № 4 (Octo 5th & Jan 13th. and Remarks pag 8 single (Jan 13th last page) cespitose (remarks pag 8) — Accord to drawg 12 inches high 3-4 dents, 8 ribs sharp; 8 radial no central spine 3 shorter than the rest, 2 above shortest one below. about 8 stigmata — Open dry ridges.

9. Mamill micromeris

10. M applanata.

11. M. macromeris? near Mier seen only once № 3 (letter Octo 5th Jan 13th) March 29th May 29 never seen on the upper Country! see living specim! —

12. Anhalonium letter March 6 1853

Living specimens

5 Ech. setispinus very short central spine, very flat. 12 rad. spines unequal, hook hardly as long. about 8-10 rad sp. Many unequal about Conepts, central 1-2 shorter stout

13. Cer. vanbelis 3 angled — shoots slend 6-7 angled, ste. letter 13 Janry last page Remarks pag 8 & 9

Dry specimens

14 Echinocactus Schottii № 2 in and label on specimen — 8 ribs — letter Octo 5th, Jan 13th 8 ribs & stout spines —

15 E. flexispinus?, areoles brown or Wislizeni?

16. Opuntia elevata similis areoles applanatis with fruit

Remarks 11

11 cf Nopal de Castil pag 4

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

14386

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Opuntiae.

O. elipticae. Nopales (Mexican)

No 1. O. Tuna (mihi) O. occidentalis (Engelm)
Erect, large flat joints obovate-rhomboid, very spinous, usually 4 stout spines of pale straw color the lowest often reflexed. Color of the petals rich lemon tinged with brickred. Pistils and filaments rather short. Carpels ovate dark-purple invested with (spinous) prickly areolae. Pulp eatable and of a better taste than those of all the other species I ever had tasted. Much sought for by the lower class of the inhabitants.

16-20 areolae on large oval Ovary

Locality. Seabeach round San Diego. California.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

14387

No 2. O. anacantha (mihi)
O. basilaris. (Engelm)

If the details sent on should prove to be different species, it is my fault, as I did not close enough compare. The one from the dividing ridge on the western edge of the Colorado desert and the other from the Gila valley about 18 miles above its mouth, which I consider the same physiographic region show much similarity, therefor I was satisfied to take only ~~the~~ dry flowers of the latter, all I could get.

MISSOURI BOTANICAL
GEORGE ENGELMANN

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

No 3. O. patens (mihi)

Cactaceæ n. sp.
near Mohavensis ?

Erect or subprocumbent, 3-4' high, joints like those of No 1. but less fleshy, areola less spinous (sub-nullate) occasionally. from 1—4. This plant occupies sometimes very extensive patches of land, which it covers exclusively. On the backs of O. Dunn I often noticed a sort of scalelike stipules covering the ~~buds~~ newly formed areola a form which I did not see on the O. patens. Flowers much like No 1. Fruit not known to me.

BOTANICAL GARDEN
ENGELMANN PAPERS

Locality. Mountain valleys of San Pascual & Santa Isabel. California.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

violacea.

No 4. O. ~~barbata~~ (mihi) = O. phaeacantha

Erect 2–3″ high; joints aboovate not fleshy; spines 1–5, light brown, red, violet color or atrofuscous sometimes cofé brown, also the epidermis round the areola sometimes ~~red~~ tinged with the colours mentioned. Otherwise the spines are long slender often contorted somewhat recurved or deflexed.

Flower & fruit not known to me.

According to the habitus of the plant I believe it is identic with a species appearing often further below in the valley of Rio Bravo and also in the vicinity of the Sierras de las Vampiros & Diablo entirely purpurascent If I not err I sent You once the drawing of one joint upon which You it asked for which reason I had drawn the spine recurved. The originae of this sketch together with the above meant plant seem to be identic.

Locality. Southrange of Sierra de las Organos near Fort Bliss. West Texas.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

10

Fa Gyrandra E

No 5. Opuntia humilis. mihi

Small plant, procumbent; joints elliptic fleshy 1–2" diameter.

Of this plant I brought home one living joint & placed it in the care of the public gardener. I believe however it shared afterwards the fate of O. vivipara (mihi)

According to its habitus this plant is identic with the original of Sketch, which I sent to Gen in 1853 among others.

Locality. Valley of the Bravo del Norte near the Cañon. (Cañon de los Lamentos of the Mexican)

I send one dead dry small specimen.

Opuntiae.

O. tessellata E

B. cylindraceae. d. gracilipines.

No 1. O. tessellata. (mihi) O. ramosissima (Engelm)

No 2. O. ramosissima β. setaceocarpa. from the Sierra Sula, wastes towards the California Gulf. probably only a freak of nature.

MISSOURI BOTANI
GEORGE ENGELM

The above forms seem to take on the west slope of the Sierra Madre the place of O. fruticosa, which occurs here only erratic.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

14388

No 3. V. sicca. (mihi) frondescens (Engelm)

GARDEN
PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

O. (Cereus ?) Schottii

No 4. Opuntia glabra (mihi)

Truly arborescent with one stem and crown. 7–8' high. ~~Branches~~ ramification divergent – divaricate; younger branches subtuberculate last years and older branches smooth, all bright green. Areola and spines resembling those of O. fruticosens to which it seems to be near allied.

Flowers greenish – yellow shaded into brick red. Petals linear lanceolate – erected.

Skeleton almost entirely solid – wooded.

Locality. Desert heights near the Maricopa villages in the Gila valley.

Whether this species may belong to the O. grazilevis I am not certain. Its ~~stem~~ only finger thick older and younger branches point to that arrangement, whilst the bold stem 4–5" diameter may classify this plant with the O. robustiores.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

B.

3. robustiores. (Choyas of the Mexican)

Nos. Opuntia Whipplei (DC) polytuberculata. (mihi)

8–9 feet high dendroid, ramifications remote-divergent; joints often 1' long by 1" or little more diameter. Tubercles numerous more flat, densely arranged greyish green; areola ovate more or less spinous, spines greyish-yellow the longest 8–9 lines.

Inflorescence terminal; carpellum pitcher shaped frequently very much elongated as to form at the former clubshaped. Corolla red.

A regular desert form, though keeping more in the levels upon the drift soil between the sierras.

Skeleton reticulate.

Locality. Llanos between the Gila and Tucson.
Sonora.



143891

mamillata.

No 6. Opuntia mammillifera. (mihi)

Dendroid with one stemp and densily ramnificated crown 5—6' high. Joints dark sepgreen, longest 4" by 1,5" diameter, ~~acto~~ mammilla oblonge-ovate full swelling developed like a womans breast. areola ovate ~~on~~ tomentose with ± 3—6 short lightyellow spines of which the longest are 7/8" length. all the spines reflexed.

Inflorescence terminal. Petals very short hardly longer then the sepals. Corolla bright pink or purple. Bacca spinous ovate-cupshaped not or very little grooved.

Locality. Southrange ~~of~~ the sierra Baburquibari and from there South East into Sonora.

This form appears only upon better soil and where rain occurs more regular. Colon, form and habits seem to point to that end.

dendroid shape of Opuntia mammillifera mihi.

MISSOURI BOTANICAL
GEORGE ENGELMANN

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

echinocarpa

No 7. Opuntia sterilis (mihi)

N.B. O. sterilis of course is wrong I meant to name it O. desertrum.

Frutticose 4–5' high virgate or divaricate joints clavate meagre 8–10" long, very tuberculous tubercles lax somewhat compressed, areola comparatively small bristled with more or less horrid ~~spines~~ and sheathed spines, the longest of which measure 1,5".

Inflorescence lateral towards the upper ends of the joints

Carpels deeply grooved and very spinous.

Locality Western parts of Pimeria alta and the deserts on both sides of the lower Colorado.

BOTANICAL GARDEN
ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Opuntia ? Fulgida

No 8. Opuntia lux Coyotis. (mihi)

(Vela de coyotis of the Mexican)

Arborescent one rarely more stems. ramification coaretate; younger and last years ~~joints~~ very densly arranged; joints ventricose - cylindric 4 - 8" long often by 2" diameter; tubercles more or less dilated areola ~~oblong~~ ovate - circular with 5 - 8 sheathed; Spines of 12 - 15''' long. Sheaths broad strawcolored and shiny, whereby this plants appears to eyesight upon a distance of 2 - 3 miles.

It is not improbable that this gave rise to its popular name.

Inflorescence terminal. Corolla purple. Baccafruit round, proliferous often forming ~~clusters~~ pendulous clusters of 10 - 12" length.

Petals linear - or clavate - lanceolate often revolute.

Skeleton reticulate.

Locality seems chiefly to belong to the basin of the californian Gulf and the lower Colorado, occupying however there more elevated positions.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

O. fulgida

Thurnbeigilia
O. D. Johnson

12

Remark to O. lux Coyotis. (mihi)

In passing over the western borders of the Colorado desert I noticed a form of a Opuntia (cylindrica opuntia) not seen before by myself. One main stem bore a large number of joints branching out laterally in a somewhat deflexed direction. These not of very great length gave the plant an appearance like a Sphagnum among the mosses. The ferruginous color of its very numerous spines and consequently of the whole plant added so much to the similarity that I was tempted to name it O. Sphagnoides. Discovering however no essential characters by which to distinguish it from O. No 8. I concluded it to be only a younger half grown form.

Areola nearly circular convex with 8–9 spines.



Skeleton of the above, also much like that of O. lux Coyotis.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

No 9. Opuntia arborescens. (Engelm)

This plant well known already deserves perhaps its specific name less then several others of the Onoya tribe.
It occurs rarely with one stem and appeared at least to me almost invariably fruticose more or less erect.
A more distinct character of its habitus is the nearly strict verticillate arrangement under which are its horizontal or pendulous joints, branching out from the main joints of its stems.

No 10. of the Cacti in Emorys / th report may partly point to the above species, the detail drawing however shows more the habitus of my O. polyhuberculata with its elongated clubshaped bases.

N.B. I do not recollect of having ever seen this species on the westside of sierra Madre.

MISSOURI BOTANICAL GARDEN
GEORGE E.
copyright reserved

14390

No 10. O. prolifera corallina. (mihi)

Boldest form of all the Opuntias, 8–10 feet high, frutiçose-divaricate much spreading often occupying much space that is more then 200 □ feet. Ramification especially above very dense irregular, joints very numerous from orate-to ventricose-cylindric 5–6" + 2." Epidermis dark but rich green.

Inflorescence of which I marked only undeveloped buds in November terminal; corolla purplish.

Skeleton reticulate, dense woody. The meshes corresponding to the tubercles of the living plant become by age gradually closed, which, as it appeared to me causes finally the dying of the plant.

The large bushes round San Diego and its vicinity appeared to me often like coral-reefs of the atmospheric ocean also thus unapproachable, therefore the name I fixed to it. field

Areola, spines no essential characteristics, as they change very much in number and arrangement upon the areola. Especially are the upper ones often absent.

BOTANICAL GARDEN ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10 cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Serpentine E

No 11. Opuntia littoralis (mihi)

4–5' high

Bush erect with a more slender ramification than the former sometimes procumbent. Joints often 1' long by 1" diameter and mostly diverging from the main stem under an ~~ans~~ acute angle.

Tubercles well developed, areola and spines resembling those of No 10. The latter pale yellow or rusty brown.

One single late blossom was procured, the corolla of which is variegated with lemon yellow and purple, bacca cupshaped, grooved and spinous.

This plant being the first of the cylindric Opuntias coming from the Atlantic side I named

O. ~~latissima~~ littoralis.

Inflorescence terminal and lateral.

If there is any prudence to refer to the bad illustrations of Cacti in Emory's former Report No 11. O. californica might be looked at under my No 11.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

13

C.

O. spuntiae humifusae.

No 12. O. spuntiae glaucae. (mihi) Emory E. Jr. del.

Low and apparently leading a more terrestrial life no ramifications, the joints proliferous, from 12–18" high. Areola concave tomentose bristling with sharp spines of a pale straw color inclining to shade a little into purple, becoming older however they all adopt a brownish grey which together with the dull grey green color of the epidermis gives the whole plant a dead-like grey appearance, hence the name.

Spines 10–12 fasciculate, erect 12–15''' long. Inflorescence terminal. Bracea dry, oblong-ovate very spinous. Corolla sulphure colored tinged with purple. Seletion woody subreticulate Ovary [illegible].

Blooms in August & September.

Locality. Desert of the Colorado occupying there extent patches of loose drift soil. Under the protection of this plant rabbits and other mining quadrupeds have their dwelling places, which thus form regular fortification against any intruder except snakes.

a O. purpureo-spina (mihi) which I found first in, similar locality, I conclude now, must be identical with the above.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

14391A

I deposited in the above sheets drawn up for Yourself the copies of fieldnotes and sketches. It was an impossibility to send You the originals, because they are all embodied in my fieldjournals together with other notes.

Posterior notes, conclusions of an eyewitness and especially more elaborate sketches or drawings, exhibiting the habitus of the plants in question, especially the larger ones at C. gigantea, Thurberi and Sinu or the larger Opuntiae You must let me have more time. In my views on scenery, which shall illustrate the maps of the Boundaryline I will try to give those singular landscape-features of the country where those large cacti occupy a prominent position.

In regard to Stanly's original drawings, I was not able to procure them. God knows where they are and whether they would be of much use.

The label to Mammillaria dasyacantha was not misplaced, it belongs to the sp. with yellow recurved spines.

M. Scheeri is not known to me.

In regard to the specific names given in the field by myself, they have of course to be left out, wherever priority on the one or impropriety on the other side should command it. If You mean to honor any of our cooperators in Cactus matters with Godfathership

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

14391B

I shall heartily consent to it. Only if You should find it practicable, You will have to let open one to be named by myself, that is I wish to do what Yourself would not be able to perform.

~~Yet~~ With the above I close for this time, thus I am

Sincerely Yours,

Arthur Schott

Maj: W. H. Emory is actual Major in one of the Cavallery regiments.

BOTANICAL GARDEN
GELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN


0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI
BOTANICAL
GARDEN

A Schotts Notes 1856

1

Echinocacti.

Nigger heads, Turkshead (Texans); Mexican Turkshead)

Bisnagas Viznagas (Mexican.)

No 1. E. Stella maris.

2. E. cottonifera = Engelmannii

3. E. caput Medusae. mihi = Emoryi

4. E. Davis Herculis, = Le Conteii

5. E. Wislizenii. ?

6. E. Catechrostatus

7. E. Lecontei

Echinocacti.

No 1. Echinocactus Stella maris. mihi.

E. viridescens. Nutt.



Areola.

Ribs. 13 – 21 ribbed. 4 central spines annulated somewhat ~~recurved~~ recurved

This plant is usually single growing, sometimes are 3 – 4 associated together.

San Diego Seabeach.

Vertical sections of the brown & persistent flower

V.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELM

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

14392

No 2. E. cottoniferus. mihi.

E. Engelmannii v. E. polycephalus E & B



Bacca and Areola of E. cottoniferus.

caespitose; heads 10–15" long by 5–8" diameter

~~ribs~~ ribs 10 — 15; Younger specimen showing but 10.

Areola round wooly; central spine longest 2" and more, radial but 1" hard but brittle; spines annulate subescent; carpels wooly elongate-ovate. Flowers persistent. Inflorescence irregularly arranged on the apex, Siluvine banks of the Gila & Colorado.

Sometimes in bunches of 5–7 heads



MISSOURI BOTANICAL GARDEN
ENGELMANN PAPERS

V.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Emoryi

No 3. Echinocactus Caput Medusae. mihi.

E. Emoryi. Engelm.

Gigantic, 19–20 ribbed. Ribbs straight or oblique,
greyish green. Largest plant 2,5 ' by 20–22 " diameter



Areola.

Transverse section of the centrespine.



Apex with the inflorescence.

Cerro Medio, Pinto
Sierra del Pinacate, S. de la Union
S. de Sonoyta.

Flowers in Aug. & Sept.
Petals atropurpureous.

V.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Cactaceae. Echinocacti.

(2)

E. Leconteï

No 4. Echinocactus clavis Hercules. mihi.



3' high by 10–12" diameter. Inflorescence somewhat resembling that of E. caput Medusae but the flowers are ~~solitary & less dense~~ arranged more scattered.

Areolas ovate or round, 2" distant. Ribbs crenate;

Blooming Aug. & Sept. like E. caput Medusae. Flowers somewhat bell shaped, petals lemon-colors with a brownish hue along the nerves.

Locality. Gila and Western parts of Pimeria alta. I found them in greatest abundance on the slopes of the Sierrita de Sonoyita, where they made a show by being associated together to 5 and 10 individuals and appearing like young Cereus giganteus.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

No 5. Echinocactus novogalensis (mihi).
(sketch natural size)

10 ribbed.



10 ribbed

Echinocereus ??

MISSOURI BOTANICAL
GEORGE ENGELMANN

Whether the above plant be really an Echinocactus I am not certain. Only two individuals of it were procured as living specimens. Circumstances however forbid afterwards the abandonement of a whole series of living cactuses in the sierra del najarito and so I have nothing to offer but a copy of the sketch which I made on the spot, where it was growing among Mammillaria stoloniformis (mihi)

The locality of its discovery was „el nodreso" on the foot of sierra del najarito near 111° long.
32.20° lat.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

14393

No 6. Echinocactus Lloydii costatus.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

Locality. Southrange of the sierra de los organos
near Fort Bliss. West Texas.

brought here
Several heads, as living specimens, they are now in care
of the public gardener.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

7) Echinocactus Wislizeni. Engelm.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

19 Dec.
1 high.
1' diameter

Near Canyon Diablo (Pila) W western part of the Boundary Line

Flora Arizonica?

Flora Arizonica Coryphantha?

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

14394

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Echinocactus ~~capitat~~ [illegible] Wislizeni

claviceps Hercules.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

14395

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

A. Schott's Remarks

Jan 1

Nopal, the characteristic plant of the flora of Mexico and therefor placed as an emblem in the arms of this empire. The genus Opuntia ~~to~~ which the Nopal plant (Prickly pear) is belonging to, is in the regions along the Rio Bravo variously represented by numerous species. Most of them are ranging from the Seacoast up ~~to~~ through the whole country. Of the various species, which have come to the knowledge of the travellers and collectors several may be considered only as subspecies or mere varieties. Like the Mesquite wood or Yucca among the exo- and endogenous plants of the country takes the Nopal-plant the most prominent station among all the Cacti of Texas. Its ~~existence~~ existence on the soil of this country is of the highest value and undeniable importance, though it is usually looked at by travellers, ~~as~~ especially by American and other foreigners as a "course upon the land", created only to sting and ~~to~~ to occupy valuable space. What however the speculative Anglo American, Dutch or German has not yet properly recognized, the contemplative Mexican, half Spaniard half Indian, became fully aware of a long time ago. The usefulness of the Nopal is well appreciated in Mexico ~~thus~~ and also in Western Texas.

In the first place the ~~organism~~ organism of the plant answers perfectly the purposes of nature. ~~As~~ Occupying immense tracts of ~~land~~ dry arid land, where hardly any other plant is able to prosper, the Nopalplant possesses a strong power to absorb atmospheric moisture in regions which in this respect

Remarks on Texan plants by Arthur Schott

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

seem to be forsaken.

The Nopal produces on the waterless sterile surface of the desolate regions of Western Texas precious food thus well for the animals of the wilderness as also for those, which aid and accompany man in his domestic endeavorings of civilization. It is for the latter reason, that the Mexican considered in former times the Western parts of Texas as the most valuable grazing country and in fact before its prosperity was so thoroughly annihilated by the hosts of the red man hundred thousands of heads of cattle, horses, mules, sheep and goats ~~and~~ have been raised and sold annually from those parts of Texas. A proof of this fact is still alive in those ~~flocks~~ herds of innumerable horses, mules and cattle, roving even at this moment in a perfectly wild state through those regions already mentioned.

The young unarmed shoots of the Nopales offer, as it is already stated, valuable food to all animals living on vegetables at a time, where the whole store of other vegetable food of the past season is consumed and grass not yet out again. This is during the months of January and February and also during the season, when all other plants in the country are dead asleep under the influence of the scourging midsummer sun. For the domestic animals the Nopal food can be prepared at every season, as the Mexican do by parching off the prickles by means of an open fire.

The people of Texas and Mexico use pieces of Nopal split

in two for covering and protecting wounds and bruises of animals and even men. I myself had often opportunity to observe ~~a knowledge~~ good success of such an application.

Another peculiarity of such split pieces of Nopales is the effect of clearing muddy water in time of some few minutes ~~muddy water~~. After the prickles being parched off such a piece is fastened on the end of a small stick and thus used ~~it~~ to stir ~~the~~ up the water to be cleared. The earthy matter by this proceeding forms a sort of as floccly sediment, which after a short time falls rapidly to the bottom.

The fruits of the Nopal even of the common are eatable, though they seem to be not entirely indifferent. I saw several times persons eating 5 – 6 pieces, without experiencing any inconvenience, whilst I saw at another time a young Mexican getting a pretty severe chill * after having eaten ~~the half~~ only one half

Of Nopales even among those resembling most each other in appearance and habitus several species seem to occur in those regions along the Rio Bravo. It was in consequence of adverse circumstances, that I was deprived of the opportunity of studying the various members of this genus more close in their home. I think however there is some slight difference in the outer appearance of this plant ranging down on the whole line along the Rio Bravo, which I have travelled over. Thus I met with the Cochenilla not before the summer 1853

MISSOURI BOTANICAL GARDEN
ANN PAPERS
copyright reserved

4

during which time I was engaged to survey from the town of ~~Laredo~~ Laredo down to the vicinity of Camargo. Whether it was an uncommon rainy season favoring the growth of the precious Cochenilla, ~~is~~ or is this tract of land really the home of this insect, I can not ascertain. Higher up as Laredo I do not remember of having seen the Cochenilla. The appearance of it on the Nopalplant is much like mildew. The Mexican gather it frequently here for their own domestic use. After smashing it to a perfect dow a flat cake is formed of, which, exposed to the rays of a subtropic sun becomes soon perfectly dry and hard. In this shape it can be preserved for a long time. According to my knowledge the Cochenilla of this part of the country is not yet brought to any commercial account. ~~in this manner~~. An other species though not growing wild ~~in~~ sometimes is met with in the country on the Rio Bravo. The Nopal de Castil, as it is called, seems to have been brought by the Mexican from the South. Thus it is to be found here sometimes under the protection of and their inhabitants Ranchos. I never had opportunity to see blossoms or fruits of this plant, the branches of which are more elliptic round and full, but not so large as of the common one. They are also more whitish or pale green. The fruits, as I was told, are more sweet and entirely harmless, the plant generally more gorgeously growing. Thus I saw a specimen in the Mexican town of Santa Rosa twenty five feet high and occupying a space of more than 1414 squarefeet. The young sprouts of an Opuntia usually dressed in different style and eaten in Mexico probably origine from this species. ———

Opuntia fruticosa.

Tunita, dimonutive form of Tuna is the fruit of Opuntia fruticosa, which is very common in every locality, where other Opuntia's have their home. Most of them are ranging through the whole of the Lowlands, even down to the shore of the saltwaters. The fruits being of a sourish taste are ~~eatable~~ eatable. In answer to remove those hairlike prickels peculiar to them the Mexican place a portion of the tuna's in a vessel with some water, brushing and cleaning them by means of a broom.

Like the nopales the bearers of the tunita, which might be called, nopalito's have also only some few hours during the ~~middle~~ noon of the day, when they open their blossoms.

As the whole family of these plants is armed by nature it is frequently used for ~~building~~ planting living fences. The easy manner by which ~~they~~ even the smallest cuts are taking root favor the application for such purposes very much.

Arthur Schott Remarks on Texian plants.

14397

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Coyonoste, is the Mexican name for the Opuntia arboresc-
ens. This plant is growing only in the upper country.
I met it ~~for~~ first on the ElPaso road between Leon-
springs and the first striking of the usually dry river
„Limpia." It is on this line, where the ~~Limpia~~ strata
of secondary limestone and even of cretaceous rocks
appear as having been protruded by the tops of
volcanic rocks. On these, which I may call igneous
rock Islands, I saw the Coyonoste first, almost ~~uninter-~~
regularly
~~ruptedly~~ appearing and disappearing with them, until re-
maining constantly in sight after reaching the Sierra Diablo
which is towering up right close on the before mention-
ed Limpia river

I never could learn the meaning of the word Coyo-
noste, but I was informed that the inhabitants of
the country, Mexican and Indian, use the sap of this
plant as an addition for brightening up the dye
of the Cochenilla.

Opuntia arborescens

Remarks on Texas plants by Arthur Schott

14398

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Bisnaga or Visnaga under this Mexican name a small Mamillaria was shown to me by a gentleman, who was living for a ~~long~~ long time in Mexico. As he had added the remark, that this plant is common in use to prepare preserves of, I suspect that there must be some mistake about the identity of the plant. I rather believe that the name Bisnage is applied to those Cactuses, which are commonly called by the Texians Turks- or Nigger heads and which bear the generic name Echinocactus. May be that only E. Wislizenii ~~and~~ E. Texensis are meant with ↑ and E. Visnago of these plants the fleshy inside is eatable, as I often tried it myself. The taste resembles much like that of fine turnipes.

The Mexican confectioners frequently cover pieces of this eatable with a sweet coat of candy, which proves to be indeed a fine sweetmeat.

Remarks on Texian plants by Arthur Schott

14399

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Organos. With this name the Mexican on the lower Rio Bravo design a Cactus of the family Cereus. Thus far as my knowledge goes, two or ~~three~~ different species are understood by this name. ~~Throughout the~~ Throughout the cretaceous formation there is one growing cespitose, in dense clusters. Its octogonal branches are somewhat falcate with an obtuse appex. Thus the whole plant has the appearance of a bundle of upright standing cucumbers. For this reason the Texian of the english tongue call it „ Cucumber-plant, whilst the Mexican call it „ Organos " (organpipes). Its eatable fruits offer to travellers one of the finest refreshments during the months of Mai & June. Shape and taste of the ~~Pitallas~~ Pitallas, as these fruits are called are much like large gooseberries and even the inside part, with its black seeds imbedded in a whitish jelly resembles much to the latter. Higher up in the Valley of the Rio Bravo, where a more solid limestone is taking the place of the softer cretaceous Greensand an other Pitallaplant occurs among the former, much like the former though larger in all its dimensions. The fruit of this is also eatable but more sour and has therefor the name „ Pitalla acre ", whilst the plant itself is also called Organos.

Further below where the cretaceous rocks give room to

C. variabilis diluvial and alluvial deposites an other Organosplant is making its appearance, comprehending two species, one tetrangular ~~the~~ an other triangular. The growth of both

Remarks on Texian plants by Arthurs Schott

is not cespitose. As I was informed by the inhabitants of those regions the blossoms of these organos are yellowish white; about their fruits I could not learn anything. Whilst both the former prefer open dry ridges, the latter like to stand more under the protection of other plants in more fertile soil.

In other parts of the country the name „Los organos" seem to be applied even on quite different plants, if they have only a somewhat like appearance.

Thus for instance I heared mention the Bonpuiera splendens under the name „organillos."

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Cereus.

No1. Cereus giganteus. (Engelm) Sahuaro (of the Sonorians)
Except sketches exhibiting habitus and deviations of its growth generally known nothing has been collected towards scientific description.

In answer to Your questions:
I never could learn anything as how old they may grow and the irregularity of the seasons and the hygrometrical conditions of the air do, I believe, not allow to any conclusions as to their annual growth. In fact for the there exists no regular fiscal- or solar year for the vegetable kingdom in the western wastes of Chihuahua and Sonora and even Texas.

Every rain actually falling causes a new spring sometimes such occur during certain months quite frequently sometimes these countries suffer drought for 16- 18 months.

Cer. giganteus
with flowers

Drawings to show the naturelike appearance of the plant in question together with skeleton and irregularities I want more leisure for than I am now permitted to expend.
The present is all I possess on fieldnote or sketches. A drawing of the fruit I would have to make from memory aided by material of dry specimens.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

JOHH1

No 2. Cereus Thurberi (Engelm)
Pitahaya of the Sonorians.

10–15' high. fruticose diving 10–15 yellowish-green ascendent stems from one root. Stems articulate, lower joints 2–3 upper 5–6 feet long. diameter 4–5". Ribbs 13–14 very little relief

Areolas distant 12–15''' tomentose when young ~~set~~ bearing 7–9 or more slender setaceous atrophus short spines, the longest of which are 6–9''' long.

Orange
Bacca globose, olivecolored, size of a small ~~orange~~ Shell soft and like all the eatable fruits of the genus invested with numerous tomentose areolas and bristle like thorns, which however give way easily in their state of maturity. Wool of the areola dirty white.

Flesh of the bacca bright carmesin richly imbedded with numberless black shiny seeds.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

Skeleton forming a weak woody ring consisting of flattened pieces of fibrebundles. The tube of the dead skeleton loosely filled with a dark brown dry powder, the remains of a loose cellular tissue.

piece of skeleton natural size.

Transverse

White
Flowers ~~of a white~~; petals ~~and sepals~~ olive colored turning gradually into a dirty white, where they are to be replaced by the petals. ~~approach the petals.~~

Inflorescence not confined to the vertex of the stems but often running down 1–1½ below it.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

No 3. Cereus Sina. (mihi)
Sina, Sinita of the Sonorans.

Pilocereus Schottii
? Sina
? pachystichus
? crassipiles

Fruticose, 8 — ~~20~~ 10 & more stems ~~often~~ stems on one root; often several plants associate together occupying in dense clusters whole patches of land. 2 — 4 joints forming one branch 8 — 10 ' high; diameter 4 — 5 ". Older joints 4 — 6 — 7 gonal Areolas tomentose when young bearing a dirty colored wool and pendulous bundles of slender bristle-like greyish-white spines of 1,5 — 2 " length. Older areolas naked and showing 4 — 6 radial & 1 central spines all ~~be~~ stout and not over 1 — 4 lines long. These spines however seem to lose themselves afterwards, leaving the stems of this plant entirely unarmed.

Inflorescence like ~~#~~ C. No 2. Flowers somewhat hidden under the long bearded areolas of the upper joints persistent

Sepals and lower petals olive brown, petals dirty flesh-color. Corolla tubular like that of C. Giganteus. Ballday, 3 — 4 ''' diameter in length, bright scarlet, seeds bright-black.

Locality. First seen on a small sierra near Sonoyta and from there South East towards Sta Magdalena and other places in that section of Sonora.
No 2 & 3, often occur together.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

14402

No 4. Cereus Hystrix (mihi)
C. Engelmannii. (Parry)

Pitahayita or Pitahaya di suelo (Sonorians.)

This plant seems to be known already.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

Locality. Common in the Sila valley especially near the casa blanca above the Pimo villages.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

8

No 5. Cereus Senecipart. (mihi) pectinatus var rigidior

Popular names the same as with Mammillaria arachnidea. (mihi)

(6–8" high 2" diameter)

Cylindric / towards the apex a little attenuate ~~verde~~ apex applanate with an excavate centre.

Ribs 20–22. Areola linear-lanceolate naked.

Spines radiate, strictly horizontal horny, stout, yellowish-purpurescent transparent 14–16.

Inflorescence on the outer margin of the perigonium. Sepals to the lip invested with bundles of delicate white spines, petals bright pink or rosecolor and along the nerves of a darker hue.

Bacca globular / dark green often 0,5–1" diameter afterwards in the state of maturity turning brownish. Pulp acidly like that of several others of the genus. The flesh of the plant is also eaten by the inhabitants of the country. Rest fibrous.

Locality. Pimeria alta in the Sierras. First seen in the valleys and arroyos of Sierra del Pajarito.

The color of the spines tinted in red, yellowish and white often forms rings around the cylindric plants as if by their different hue indicating some thing like annual growth

Obs. Rem. What I suppose to be Cereus caespitosus has an ovate circular areola with similar and alike arranged radiate spines like C. Senecipart, but on the latter I never noticed a central spine. Habitus of the former is also different.

0 1 2 3 4 5 6 7 8 9 10 cm
copyright reserved
MISSOURI BOTANICAL GARDEN

9

Dasyacanthus 2

6. Cereus crassus. (mihi)

Subcylindrical, ventricose ~~obpyriform~~. ~~naked~~, ~~terete~~ cylindrical, ~~acuminate~~ canose or subcespitose upper conic; 17–18 ribbs; areola ovate, spines fasciculate covering the whole areola, lower longest ~~to~~ 3–4 ''' long, white and reddish or pink annulated and tipped, base ventricose ~~inflated~~ inflated;

Flowers or fruits not known to me.

I send one living specimen of C. crassus (mihi)

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

Locality South range of Sierra de los organos near Fort Bliss. West – Texas. and farther below in the valley of Rio Bravo.

Remark. No 11. Cereus rigidispinus (mihi.) In habitus almost is the same entirely alike to No 6. has for me no other marks of distinction between both except the number of spines, of which my C. rigidispinus has shows only half the number of those C. crassus has. Number of spines not being essential enough I shall not be surprized to see both those Cereuses ~~was~~ members of the same species

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

~~6~~ Cereus senecionis. Mr Monkendel
grey headed. cabezito de viejo. Cylindrical
~~erect~~ ~~a~~ upright ~~spreading~~ 20-22 ribbed
~~green~~ dirty-lanceolate ~~glabr~~ naked.
areolas ~~close~~ very close
~~and spines all~~ but radiate, ~~yellow~~ horny and
short & horny, yellowish-purpurescent
~~brownish~~ translucent. Rests to 6-8" high.
Flowers large pink or bright rose color
~~crown~~ ~~crowning~~ appearing on the periphery of the
apex. like Bacca ~~globose 1½-1" elastic~~
globose 1½-1" elastic
bacca invested with bundles of small white
spines, which and in the state of maturity de-
a dorus spines. Bacca glabrose often of 0,5-1½
diameter. ~~Fruit~~ Bacca like the Bithaya
eatable, ~~also~~ The flesh of the plant is also
eaten by the inhabitants of the country,
Primeria alta. Bot Informs

Spines resembling those of Cereus diverse
arranged
but straight and horizontally around the
areola, the points ~~crossing~~ overreaching
each other.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

14403

Cereus horridus.
20 – 2 ½ Ins. –
Mrs C. Seneciput. mihi.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS



12-16-17.
Spines



areola of
C. cespitosus Engelm.?)

Rose or pink color.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Cereus albispinus. (mihi)

Plant cespitose, dark green spines short white.

13 ribbed, cespitose heads

obtuse conic apices excavate.



Areola.

1

Head of rio San Pedro.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

14404

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

caespitosus E

7

No 8 Cereus dipsacus. (mihi)

C. caespitosus of Dr. Bigelows living collection in the public gardens ~~here~~ at Washington DC.

Cylindric, apex obtuse, 13 ribbed; ribs little interrupted areolas. 2‴ distant; areola ovate-lanceolate; spines all radial 16–18 ~~spines~~ or more spines, adscendent-recurved, lower spines longest to 3.‴ upper shortest. Sp. horny yellow and tipped tinged with purple

. Root fibrous.

areolae.

Flowers and fruit not known to me.

Locality between Escondido spring and Pecos river. West Texas.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

14403

No 9. Cereus pitahayita. (mihi)

A. Cespitose and in habitus, number of ribs, ~~areol~~ areola and spines almost identic with the species on the lower Rio Bravo, known as Cucumberplant, for which I ~~took~~ took it in the remote locality of the Sierra del Pajarito (Sonora) Seeds, with the dry brcew~~n~~ and persistent red flowers are at hand. Part of them was transmitted to You. I think the difference between these different species would be enough to pronounce the Cereus pitahayita a new species, be it with this or an other name. To my C. hysterix the C. pitahayita bears ~~no~~ but a general resemblance whilst even a more practized eye will easily mistake it for the cucumberplant.

I am sorry to say, that a small living species ~~being~~ ~~place~~ arriving here must have been lost under the hands of one of the hands in the public gardens.

MISSOURI BOTANI
GEORGE ENGELMA

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

?

No 10. Cereus decacostatus. (nichs)
subcespitose
Conic + cylindric; 10 ribbs; cespitose or nearly so.



flower & fruit not known to me.

I believe it is one of those Pitahayas bearing cerei, the pulp of their buds is more sower and therefore called by the Mexican of the Rio Bravo valley " Pitahaya agre."

I brought home one living species now in care of the public gardener measuring 12" high & 4" diameter.

RI BOTANICAL GARDEN
ENGELMANN PAPERS

Locality Southrange of sierra de los organos, near Fort Bliss. West-Texas.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

(cespitosus ξ

12. Cereus albispinus. (mihi)

cespitose

Obtuse-conic; 1) length and diameter almost alike; 13 ribbs; ~~dark green~~ areolæ 3‴ or more distant; ribbs deeply interrupted; areola linear-lanceolate or ovate lanceolate spines short, ~~wart and~~ the lateral ones being longest from 1—1,5‴ pure white and harmless.

The shape of the areola with the spines reminded me as to name this species „pedicularis!"

No flowers or fruit known to me.



Locality. Head of rio San Pedro in West-Texas.

1) The living specimen here ~~have~~ are nearly 2+2!

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

1. Mammillaria tenuispina (mihi)
= M. Grahamii. Engelm. already known.

mam tenuispina = M. Grahamii
petens = " "
ambanda = M. recurvispina n. sp
robustispina ! n. sp
nida Dibdi = M. macromeris
Hrobilisformis = M. radiosa
acicularis = M tuberculata
pygmaea = M gummifera ?
longimamillosa = M radiosa ?

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

recurvispina E

2. Mammillaria arachnidea (mihi)

Cabezito del viejo, (mexic) Greyheadlet, Oldmanshead, (engl.) Graukopf = oder Grauer Kopfgitter (german)

Globose or subglobose, apex somewhat applanate 6-8" diameter, mamillas ~~somewhat~~ more or less compressed densly arranged and covering each other like scales. (subsquamose) Areolas ~~ovate~~ oblong & tomentose when young. Spines, one central 6-8''' long radials 12-14; ~~as all the rest~~ like the former contorted recurved, white & brown towards the point; transparent when young. central spine 6-8''' long radial 2-3'''. Suture naked and perceptible to the base of the mamilla. Flowers, somewhat campanulate irregularly distributed over the apex with an inclination to appear towards the perigonium. petals bright lemonyellow darker in with a brownish tint along the midrib.

Flesh like that of Cereus senilis mihi (mihi) eatable and used by the inhabitants of the country.

MISSOURI BOTANICA
GEORGE ENGELMA

Locality. Eastern parts of Pimeria alta especially in the sierra del pajarito.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

14406

3. Mammillaria patens. (mihi)
= M. Johanni C

Growth is low spreading bunches, therefrom the name I gave.

BOTANICAL GARDEN
ENGELMANN PAPERS

Of this plant but one dead dry specimen was found in the Gila valley near the Pimo villages.

I send part of it to look at.

In the public gardens here I saw some days ago a living specimen of this much resembling ~~apparently~~ other species.

No flower no fruit.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Valdivia

4. Mammillaria robustispinaa. (mihi)

Growth cespitose; mammillas distant one inch long or nearly so; sulcus very much marked and densily tomentose like the areola.

Spines two central, upper is curved.

Radial spines of ~~light brown~~ eight brown above deeply shaded and ~~varied~~ varied by dark brown reverse side whitish. Spines ~~and~~ 1 11 long very stout especially the central on the base of nearly 1 111 diameter.

Flowers 1,5 11 long, tube very narrow, corolla campanulate. Petals bright Saffron yellow.

The only specimen I found near the sierra Babonquibari was sedmingly mutilated by his grazy llamos prairiefire, thus the parts of it do not make much show.

The unripe seed I sent under the name M. rigidissp. can not be but identical with the above.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Mammillaria nidus diaboli.

Nihi.

before M. stalagmites. (mihi)

living specimens.

Tuft consisting of
50 – 60 heads.

Locality. Cerro de los Pinos on the
other side of Buenos Aires.

A. Schott

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

14407

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Cactaceae. Mammillarieae. Mammillaria ?

4

S/ Mammillaria mirabilis diabolis. (mihi)

Growth. Hemispherical tuft of 50–60 heads, arranged very close together.; whole mass 1½ foot in diameter and 6 inches high.

Tubercles long (& taper) 3–5 central and as many radial spines, long & slender

This plant shows itself from a distance soft almost velvet-like, but being approached it looks rather unhospitable like all the other of this order. Thus I selected its name, though it might be Your M. aggregata. Engelm's Report 1848. Appendix pag. 157. & Fig. 1.

Locality. Few miles below Escondido springs one days march above leaving the Puercos valley. Western Texas.

I send one living specimen

14408

redosa var leo mexicana.

106. Mamm. Macdougalii strobiliformis. (mihi)

Corte- ~~conic~~ conic (strobiliformis) 1,5 – 2" + 3" high. Root fibrous. Tubercles distant, attenuate to 1" long. Sutira on the younger mamillas subterminal. Areola round convex wooly; central spine 1 horizontally inwards bowed ~~when~~ on younger tubercles, vertically upon the right or older, strict tendency to curve ~~recurved~~ with the other radiate spines. The latter like the former of a light ~~colour but grow~~ ~~after the~~ during the course of time a brown hair-color. Radial spines 25 – 30 slender and ~~straight~~ straight. in 2 – 3 rows the upper from 6 – 8 lines long. Flowers persistent. apparently red, placed in the centre of the apex, bacca oval greenish-yellow.

Locality. El Pedrero near 111° Longit. 32.20° Latitude on the northern slope of Sierra del pajarito.

In habits much resembling to M. acicularis (mihi) the areola of which is however covered with smaller more slender bristle like pure white spines, which are arranged more fasciculate. Mamillas of this are also arranged compact, inflorescence not confined to the centre of the apex but appearing more towards the perigonium.

M. strobiliformis (mihi) shows its longest spines on the upper part of the areola whilst M. acicularis has them on both sides

I send one living specimen.

MISSOUTH BOT GEORGE ENG

0 1 2 3 4 5 6 7 8 9 10 cm copyright reserved MISSOURI BOTANICAL GARDEN

7) Mammillaria longimamma DC. var longimammillifera (mihi)

Of this species I met only one specimen near the Camp at Howards ranch (San Antonio – El Paso road). I tried to secure it as a live specimen but ~~must~~ must have lost it again.

I enclose herewith a sketch of it copied from the one ~~in~~ made on the spot.

Mam mills nearly 1" long not very compact arranged, dark green.

No flower or fruit.

12 radial & 6 central sp.

ICAL GARDEN
MANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

No 8. Mammillaria tuberculata aciculare. (mihi)

Conic with a broader base then No 6 to which it bears in habitus some relationship. 1—2" high. Tubercles very closely arranged; 1—3 stout straight upright central spines 6"'—12"' long, light brown dark with a darker point. 30—40 radial spines in 2—3 rows, bristle like, white x 3—4"' long.

Inflorescence on the perigonium.

Root fibrous.

Locality. Southrange of sierra de los Organos near Fort Bliss. West-Texas. with Cereus Eyi dispenius var C. crassus (mihi)

I send one living specimen. & fruit

No 2 Mamm. Clarkii long. mamillar. Mitis.







a
Mamm. Clarkii natural size
a. outside.
b. inside.

6

central spines 6.
radial sp. 6. pairs.
all the spines slender
greyish-white.

Locality near Howardsprings. El Paso
Co. Western Texas.

Arthur Schott

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

14409

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

No 9. Mammillaria prismatica? (mihi)

sessile of 4–5" diameter.

Applanate, spines atropurpurescent; 1 central spine; 13 radial. Carpels sessile ex centro deeply imbedded between the tubercles, which are densily arranged.



Tubercles pyramidical obtuse angular.

Spines Outer or ~~inter~~ lower radial spines longest to 6-8"' long; interior or upper 1 spines 2–3"' long. central spines short 2–3"'.

Bases bright crimson color.

Locality. Sierra del Rajarito. Sonora. near 111° long. 32.20° lat.

The living specimens procured had to be abandoned again by circumstances.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

14410

10. Mammillaria excavata. (mihi)

In habitus near allied to No 9. no 9. Size, shape of tubercles being very much alike. The centre of M. excavata however is deeply excavated forming a funnelshaped groove of 1" depth.

Flower and fruits unknown to me.

Locality with No 9.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

Living specimens procured had to share the fate with those of no 9.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN